arXiv:1211.2952v1 [math.DS] 13 Nov 2012

Pseudo-Orbits, Stationary Measures and Metastability

H. Hu, W. Bahsoun and S. Vaienti

Abstract We study random perturbations of multidimensional piecewise expanding maps. We characterize absolutely continuous stationary measures (acsm) of randomly perturbed dynamical systems in terms of pseudo-orbits linking the ergodic components of absolutely invariant measures (acim) of the unperturbed system. We focus on those components, called *least-elements*, which attract pseudo-orbits. We show that each least element admits a neighbourhood which supports exactly one ergodic acsm of the random system. We use this result to identify random perturbations that exhibit a metastable behavior.

1 Introduction

In this chapter we study statistical aspects of random perturbations of a multidimensional piecewise expanding map[1] $T : M \to M$, $M \subset R^q$. A random orbit $\{x_n^\varepsilon\}_n$ is a random process where, for all n, x_{n+1}^ε is a random variable whose possible values are obtained in an ε-neighbourhood of Tx_n^ε according to a transition proba-

---

H. Hu
Mathematics Department, Michigan State University, East Lansing, MI 48824, USA, e-mail: `hu@math.msu.edu`

W. Bahsoun
Department of Mathematical Sciences, Loughborough University e-mail: `W.Bahsoun@lboro.ac.uk`

S. Vaienti
UMR-7332 Centre de Physique Théorique, CNRS, Université d'Aix-Marseille et Université de Toulon, and FRUMAM, Fédération de Recherche des Unités de Mathḿatiques de Marseille; address: CPT, Luminy Case 907, F-13288 Marseille Cedex 9 e-mail: `vaienti@cpt.univ-mrs.fr`

[1] See [10] and [16] for recent results and for an exhaustive list of references on deterministic multidimensional expanding maps.

bility $P_\varepsilon(x_n,.)$. We consider the case where $P_\varepsilon(x_n,.)$ is absolutely continuous with respect to Lebesgue measure m. Denoting the density of the transition probability by $p_\varepsilon(x,.)$, we define a perturbed transfer operator , $\mathscr{L}_\varepsilon$, by:

$$\mathscr{L}_\varepsilon f(x) = \int_M p_\varepsilon(y,x)f(y)dy$$

where $f \in L^1_m(M)$. We focus on non-invertible dynamical systems whose transfer operators, perturbed and non-perturbed[2], satisfy a uniform Lasota-Yorke inequality. Among other things, such an inequality implies the existence of a finite number of ergodic acim for the initial system T, and a finite number of ergodic acsm for the random system.

We then define an equivalence relation between the ergodic components of acim of T using *pseudo-orbits*. Using this equivalence relation, we consequently introduce equivalence classes of ergodic acim. Among the latter, we identify those which attract pseudo-orbits and call them *least elements*. We show that each least element admits a neighbourhood which supports exactly one ergodic acsm of the random system. We use this result to identify random perturbations that exhibit a *metastable behavior.* Such a phenomena has recently been a very active topic of research in both ergodic theory [2, 3, 7, 8, 9, 13] and applied dynamical systems [6, 18].

Section 2 contains the setup of the problem, our assumptions, the notion of a least element and the statement of our main result (Theorem 1). Section 3 contains the proof of Theorem 1. In section 4 we use the results of the previous sections to identify random systems which exhibit a metastable behavior. In section 5 we apply our results to random transformations, in particular, we provide one-dimensional examples to illustrate the results of sections 2 and 4.

2 Setup and statement of the main result

2.1 The initial system

Let $M \subset R^q$ be compact[3] with $\overline{intM} = M$. We denote by d the Euclidian distance on R^q. Let $(M, \mathscr{A}, m)$ be the measure space, where $\mathscr{A}$ is Borel σ-algebra, and m is the normalized Lebesgue measure on M; i.e, $m(M) = 1$.
Let $T : M \to M$ be a measurable map, $\mathscr{D}_T$ be the set of discontinuities of T. We assume that $m(\mathscr{D}_T) = 0$, and T is non-singular with respect to m. The transfer operator (Perron-Frobenius) [4] associated with T, $\mathscr{L} : L^1_m \hookrightarrow L^1_m$ is defined by duality: for $g_1 \in L^1_m$ and $g_2 \in L^\infty_m$

---

[2] In this chapter, the non-perturbed operator, $\mathscr{L}$, is the traditional transfer operator (Perron-Frobenius) associated with the map T [4].

[3] All our results can be carried to the case where M is a compact Riemannian manifold.

$$\int_M g_1 g_2 \circ T dm = \int_M \mathscr{L} g_1 g_2 dm.$$

2.2 The perturbed system

We perturb the map T by introducing a family of Markov chains $(\mathscr{X}_n^\varepsilon)$, $\varepsilon > 0, n \geq 0$ with state space M and transition probabilities $\{P_\varepsilon(x,.)\}_{x\in M}$; i.e, $P_\varepsilon(x,A)$ is the probability that a point x is mapped into a measurable set A. At time $n = 0$, $(\mathscr{X}_0^\varepsilon)$ can have any probability distribution. We assume that:

(P1) For all $x \in M$, $P_\varepsilon(x,.)$ is absolutely continuous with respect to the Lebesgue measure. We will denote its density by $p_\varepsilon(x,.)$. Therefore $P_\varepsilon(\mathscr{X}_{n+1}^\varepsilon \in A | \mathscr{X}_n^\varepsilon = x) = \int_A p_\varepsilon(x,y)dm(y)$.
(P2) We have: $\text{supp}P_\varepsilon(x,.) = B_\varepsilon(Tx)$ for all x.

Assumption (P2) can be weakened by supposing that $\text{supp}P_\varepsilon(x,.)$ is a subset of a slightly larger ball around Tx. Our proofs can be easily adapted to show that the results of this chapter will still hold under such a slightly weakened assumption.

2.2.1 Random orbits and stationary measures

The perturbed evolution of a state $x \in M$ will be represented by a *random orbit*:

Definition 1. A sequence $\{x_n^\varepsilon\}_{n\geq 0} \subset M$ is an ε-random orbit if each x_{n+1}^ε is a random variable whose distribution is $P_\varepsilon(x_n^\varepsilon,.)$, namely $\{x_n^\varepsilon\}_{n\geq 0}$ coincides with the Markov chains $(\mathscr{X}_n^\varepsilon)$, $\varepsilon > 0, n \geq 0$.

The counterpart of an invariant measure in the case of randomly perturbed dynamical systems is called a stationary measure:

Definition 2. A probability measures μ_ε is called a stationary measure if for any $A \in \mathscr{A}$

$$\mu_\varepsilon(A) = \int_M P_\varepsilon(x,A)d\mu_\varepsilon(x).$$

2.2.2 The transfer operator of the random system

To study Markov processes it is useful to define the transition operator $\mathscr{T}_\varepsilon$ acting on bounded real-valued measurable functions g

$$(\mathscr{T}_\varepsilon g)(x) = \int_M g(y)P_\varepsilon(x,dy).$$

Its adjoint $\mathscr{T}^*$, defined on the space $\mathscr{M}(M)$ of all finite signed measures, is given by:

$$\mathscr{T}_\varepsilon^* \mu(A) = \int_M P_\varepsilon(x,A) d\mu(x).$$

Notice that μ_ε is a stationary measure if and only if $\mathscr{T}_\varepsilon^* \mu_\varepsilon = \mu_\varepsilon$. Moreover, a stationary measure is ergodic if, for $A \in \mathscr{A}$ with $\mathscr{T}_\varepsilon 1_A = 1_A$ implies that $\mu_\varepsilon(A) = 0$ or $\mu_\varepsilon(A) = 1$. Since condition (P1) implies the absolute continuity of any stationary measure, it will be convenient to define an operator, $\mathscr{L}_\varepsilon$, acting on densities. That is to say, if μ is an absolutely continuous measure with respect to m, whose density is a function $g \in L^1_m$, then $\mathscr{T}_\varepsilon^* \mu$ is an absolutely continuous measure whose density is $\mathscr{L}_\varepsilon g$, where $\mathscr{L}_\varepsilon g$ is given by:

$$\mathscr{L}_\varepsilon g(x) = \int_M p_\varepsilon(z,x) g(z) dz. \tag{1}$$

Thus, densities of acsm are fixed points of $\mathscr{L}_\varepsilon$.

2.3 A Banach space and quasi-compactness of $\mathscr{L}$ and $\mathscr{L}_\varepsilon$

We now introduce a Banach space $\mathscr{B}(M) \subset \mathscr{L}^1_m$. We assume that

(B1) The constant function 1 belongs to $\mathscr{B}(M)$.
(B2) The set of discontinuities, $\mathscr{D}_f$, of any function $f \in \mathscr{B}(M)$ has Lebesgue measure zero.
(B3) There is a semi-norm $|.|$ on $\mathscr{B}(M)$ such that the unit ball of $\mathscr{B}(M)$ is compact in $\mathscr{L}^1_m$ with respect to the complete norm $||\cdot||_B \equiv |.| + \|.\|_1$, where $\|.\|_1$ denotes the L^1_m norm.

We also assume the transfer operators $\mathscr{L}$ and $\mathscr{L}_\varepsilon$ satisfy a uniform Lasota-Yorke inequality: there exist an $\eta \in (0,1)$ and a $D \in (0,\infty)$ such that for all $f \in \mathscr{B}(M)$ and $\varepsilon > 0$ small enough:

$$\mathscr{L} f \leq \eta\, |f| + D\, \|f\|_1\, ; \qquad (LY)$$

$$\mathscr{L}_\varepsilon f \leq \eta\, |f| + D\, \|f\|_1\, . \qquad (RLY)$$

Assumptions (LY) and (RLY) ensure the quasi-compactness of both $\mathscr{L}$ and $\mathscr{L}_\varepsilon$ [4]. In particular, among other things, (LY) implies the existence of a finite number of T-ergodic acim, and (RLY) implies the existence of a finite number of ergodic acsm for the Markov process $(\mathscr{X}_n^\varepsilon)$. More precisely, we have for the operator $\mathscr{L}$:

- The subspace of non-negative fixed points w of $\mathscr{L}$, is a convex set with a finite number of extreme points $w_1, \cdots, w_l$ with supports $\Lambda_k, k = 1, \cdots, l$. The supports $\Lambda_k, k = 1, \cdots, l$ are mutually disjoints Lebsegue a.e..
- The measures $\mu_1 = w_1 m, \cdots, \mu_l = w_l m$ are ergodic: we will call them the ergodic decomposition of any acim $\mu = hm$, $h \in \mathscr{L}^1_m$, or the extreme (ergodic) points (measures) decomposing μ.

The Lasota-Yorke inequality (RLY) of $\mathscr{L}_\varepsilon$ ensures that the random system admits finitely many ergodic acsm $\mu_1^\varepsilon = h_{\varepsilon,1}m, \dots, \mu_K^\varepsilon = h_{\varepsilon,K}m$. Note that in general the number K of extreme points for μ_ε is different from the number l of ergodic components of μ. In our setting, (see Corolary 1), we show that the number of ergodic ascm is bounded above by the number of ergodic acim.

Remark 1. We point out that for certain perturbations one can prove that (RLY) follows from (LY). See [5, 11] for precise examples.

A well known consequence of assumption (LY) and (RLY) is the following proposition

Proposition 1. *Let* $\{h_\varepsilon\}_{\varepsilon>0}$ *a family of densities of absolutely continuous stationary measures of* $(\mathscr{X}_\varepsilon^n)$*. Then any accumulation point of* $\{h_\varepsilon\}_{\varepsilon>0}$ *in the* L^1_m*-norm is a density of* T*-acim.*

2.4 Pseudo-orbits, least elements and the statement of the main result

We now introduce the notion of a pseudo orbit which will be our main tool to characterize ergodic acsm. Pseudo-orbits were previously used by Ruelle [17], followed by Kifer [14], to study attractors of randomly perturbed smooth maps.

Definition 3 (pseudo-orbit). For $\varepsilon > 0$, an ε-pseudo-orbit is a finite set $\{x_i\}_{i=0}^n \subset M$ such that $d(Tx_i, x_{i+1}) < \varepsilon$ for $i = 0 \dots n-1$.

Using pseudo-orbits, we define a partial order "$\to$" among the supports $\{\Lambda_i\}$ of T-ergodic acim by writing $\Lambda_i \to \Lambda_j$ if for any $\varepsilon > 0$ there is an ε-pseudo-orbit $\{x_i\}_{i=0}^k$ such that $x_0 \in \Lambda_i$ and $x_k \in \Lambda_j$. Then we define a relation "$\sim$" among $\{\Lambda_i\}$ by writing $\Lambda_i \sim \Lambda_j$ if both $\Lambda_i \to \Lambda_j$ and $\Lambda_j \to \Lambda_i$. By ergodicity, given a point $y \in \Lambda_i \equiv \mathrm{supp}\mu_i$ and $\varepsilon > 0$, μ_i-almost any point $x \in \Lambda_i$ will enter the ball, of positive μ_i measure, $B_\varepsilon(y)$, and therefore all the pairs $(x,y) \in \Lambda_i$ can be connected with a (finite) ε-pseudo-orbit. Hence we get an equivalence relation among those ergodic components and we define $\tilde{\Lambda}_i$ the equivalence class which contains Λ_i. We write $\tilde{\Lambda}_i \to \tilde{\Lambda}_j$ if $\Lambda_k \to \Lambda_l$ for any $\Lambda_k \in \tilde{\Lambda}_i$ and $\Lambda_l \in \tilde{\Lambda}_j$.

Definition 4. $\tilde{\Lambda}_i$ is said to be a *least element*, if there is no $\tilde{\Lambda}_j$ such that $\tilde{\Lambda}_i \to \tilde{\Lambda}_j$.

This in particular means that for all ε small enough no ε-pseudo-orbit can travel from the least element to other equivalence classes. We point out that, in general, a dynamical system may have more than one least-element[4]. We now state our first result:

Theorem 1. *Under Assumption (P1) and (P2), if (LY) and (RLY) hold for functions in a Banach space* $B(M)$ *satisfying (B1-B3), then we have:*

---

[4] This will be illustrated in Example 1.

1. *If $\tilde{\Lambda}$ is a least element, then for ε small enough there exists an open neighborhood $\mathscr{U}_\varepsilon \supset \tilde{\Lambda}$ which supports a unique ergodic acsm $\mu_\varepsilon^{\tilde{\Lambda}}$.*
2. *If $\tilde{\Lambda}$ is not a least element, then for ε small enough, $\mu_\varepsilon(\tilde{\Lambda}) = 0$ for any acsm μ_ε. Therefore, for any weak-limit of μ_ε, $\tilde{\Lambda}$ is a set of measure 0.*

Theorem 1 implies the following three corollaries:

Corollary 1. *The number of acsm of the random system $(\mathscr{X}_\varepsilon^n)$ is bounded above by the number of acim of the map T. In particular, if T has a unique acim, then $(\mathscr{X}_\varepsilon^n)$ has a unique acsm.*

Corollary 2. *Let $h_\varepsilon^{\tilde{\Lambda}} \in L_m^1$ be the density of the ergodic acsm $\mu_\varepsilon^{\tilde{\Lambda}}$. By Proposition 1, any accumulation point of the family $\{h_\varepsilon^{\tilde{\Lambda}}\}_{\varepsilon>0}$ is a convex combination of the densities of the extreme ergodic measures spanning $\tilde{\Lambda}$*[5]*. In particular, relative to the ergodic acsm supported on a neighborhood of least elements, the original system is strongly stochastically stable*[6].

Corollary 3. *The support of any ergodic acsm contains a least element.*

3 Proofs

We first prove a key lemma.

Lemma 1. *Let $f \in \mathscr{B} \equiv \mathscr{B}(M)$ and $\{x_i\}_{i=0}^N \subset M$ be an ε-pseudo-orbit such that $x_j \in M \setminus (\mathscr{D}_{\mathscr{L}_\varepsilon^j f} \cap \mathscr{D}_T)$, $0 \le j \le N$ and $\mathscr{L}_\varepsilon^i f(x_i) > 0$, for some $0 \le i < N$. Then for all $i < k \le N$ we have $\mathscr{L}_\varepsilon^k f(x_k) > 0$.*

Proof. Let $f \in \mathscr{B}$ and $\{x_i\}_{i=0}^N$, an ε-pseudo-orbit, satisfying the assumptions of the lemma. In particular, suppose that for some fixed $0 \le i < N$, $\mathscr{L}_\varepsilon^i f(x_i) > 0$. Then

$$\mathscr{L}_\varepsilon^{i+1} f(x_{i+1}) = \int_M \mathscr{L}_\varepsilon^i f(y) p_\varepsilon(y, x_{i+1})\, dy.$$

By the hypothesis (P2) we have $x_{i+1} \in B_\varepsilon(Ty) \Rightarrow p_\varepsilon(y, x_{i+1}) > 0$ and by the preceding continuity assumptions there exists $\delta > 0$ such that $y \in B_\delta(x_i) \Rightarrow \mathscr{L}_\varepsilon^i f(y) > 0$. But this δ-neighborhhod of x_i can be made smaller in such a way that when y belongs to it, $d(Ty, Tx_i) = \frac{\varepsilon - d(Tx_i, x_{i+1})}{2}$ which implies that x_{i+1} is ε-close to Ty. Therefore for all the y in this δ-neighborhood of x_i (which is of positive Lebesgue measure), the integrand above is strictly positive and this finishes the proof of the Lemma.

---

[5] Note that within the general setting of this paper, we do not claim that the values of the weights in the convex combination that determine the limiting density can be easily identified. However, for certain perturbations of one dimensional maps, using insights form open dynamical systems, such weights can be determined. See [9, 2, 3].

[6] We will say that the system (M, μ, T) is strongly stochastically stable if any L_m^1 accumulation point of the densities of the extreme ergodic absolutely continuous stationary measures $\{\mu_j^\varepsilon\}_{\varepsilon>0}$ for any $j = 1, \ldots, K$ and when ε goes to zero, is a convex combination of the densities of the absolutely continuous ergodic extreme measures of μ.

Proof (of Theorem 1). We first show that functions in $\mathscr{B}$ which are supported in an open neighborhood of a least element $\tilde{\Lambda}$ are mapped by $\mathscr{L}_\varepsilon$ to functions in $\mathscr{B}$ which are supported on the same neighborhood of $\tilde{\Lambda}$. This will allow us to get a fixed point of $\mathscr{L}_\varepsilon$ in such a neighborhood of $\tilde{\Lambda}$. We denote by $B_\varepsilon(A)$ the (open) ε-neighborhood of a set A. When A is a single point x, this definition reduces to the usual definition of a ball of center x and radius ε. We observe that even though a least element $\tilde{\Lambda}$ is forward invariant, the image of a ball of radius ε centred at a point in $\tilde{\Lambda}$ may not be necessarily contained in $\tilde{\Lambda}$. However, this ball will surely subset of the open set $\mathscr{U}_{\varepsilon,1} := B_\varepsilon(T\tilde{\Lambda})$. We define inductively a family of nested open sets $\mathscr{U}_{\varepsilon,n} := B_\varepsilon(T\mathscr{U}_{\varepsilon,n-1})$ and consider the open neighborhood $\mathscr{U}_\varepsilon$ of $\tilde{\Lambda}$ defined by $\mathscr{U}_\varepsilon := \cup_{n=1}^{\infty}\mathscr{U}_{\varepsilon,n}$. This set is clearly forward invariant under T, and its closure is disjoint, for ε small enough, from the supports of the other ergodic acim; otherwise, we can construct an ε pseud-orbit linking the least element to them.
(i) Let $\mathscr{B}(\mathscr{U}_\varepsilon) := \{f \in \mathscr{B} | f \text{ is supported on } \mathscr{U}_\varepsilon\}$. Since $\mathscr{L}_\varepsilon f(x) = \int_M p_\varepsilon(y,x) f(y) dy$, the forward invariance of $\mathscr{U}_\varepsilon$ together with $x \in B_\varepsilon(Ty)$, insure that $\mathscr{L}_\varepsilon$ leaves the Banach space $\mathscr{B}(\mathscr{U}_\varepsilon)$ invariant. Then by the spectral construction of densities, we obtain a fixed point $\mathscr{L}_\varepsilon h_\varepsilon = h_\varepsilon \in \mathscr{B}(\mathscr{U}_\varepsilon)$: the measure $\mu_\varepsilon = h_\varepsilon m$, $\int_{\mathscr{U}_\varepsilon} h_\varepsilon dm = 1$ is therefore stationary. We now prove that h_ε is the only fixed point of $\mathscr{L}_\varepsilon$ in $\mathscr{B}(\mathscr{U}_\varepsilon)$. Suppose there is another function $h'_\varepsilon \in \mathscr{B}(\mathscr{U}_\varepsilon))$ with the same property, and let us define the function $\hat{h} = \min(h_\varepsilon, h'_\varepsilon)$. Then clearly: $\min(h_\varepsilon, h'_\varepsilon) \leq h_\varepsilon$ and $\min(h_\varepsilon, h'_\varepsilon) \leq h'_\varepsilon$ and thus $\mathscr{L}_\varepsilon(\min(h_\varepsilon, h'_\varepsilon)) \leq \mathscr{L}_\varepsilon h_\varepsilon$ and $\mathscr{L}_\varepsilon(\min(h_\varepsilon, h'_\varepsilon)) \leq \mathscr{L}_\varepsilon h'_\varepsilon$ which implies $\mathscr{L}_\varepsilon(\min(h_\varepsilon, h'_\varepsilon)) \leq \min(\mathscr{L}_\varepsilon h_\varepsilon, \mathscr{L}_\varepsilon h'_\varepsilon)$. But $\min(\mathscr{L}_\varepsilon h_\varepsilon, \mathscr{L}_\varepsilon h'_\varepsilon) = \min(h_\varepsilon, h'_\varepsilon) = \hat{h}$, so $\mathscr{L}_\varepsilon \hat{h} \leq \hat{h}$ and therefore $\mathscr{L}_\varepsilon \hat{h} = \hat{h}$. Let us consider $h_\varepsilon - \hat{h}$. It is a nonnegative function and it satisfies $\mathscr{L}_\varepsilon(h_\varepsilon - \hat{h}) = h_\varepsilon - \hat{h}$. By Proposition 1 and by taking ε small enough, we insure that the supports of h_ε and h'_ε will intersect the least element in a Borel set B of positive Lebesgue measure. Starting from almost any point in this set we can attain any other point in $\mathscr{U}_\varepsilon$ with a (finite) ε-pseudo-obit [7]. Then for any $x_0 \in B$ such that $(h_\varepsilon - \hat{h})(x_0) > 0$, we can apply Lemma 1 with $f = h_\varepsilon - \hat{h}$ and by considering an ε-pseudo-orbit $\{x_i\}_{i=1}^N \subset M \setminus (\mathscr{D}_{h_\varepsilon - h} \cap \mathscr{D}_T)$ which starts from almost any $x \in B$, passes through x_0, and lands in $x_N \in \mathscr{U}_\varepsilon$ we have $\mathscr{L}_\varepsilon^k(h_\varepsilon - \hat{h})(x_k) = (h_\varepsilon - \hat{h})(x_k) > 0$, $k = 1, \cdots, N$. This is equivalent to say that $(h_\varepsilon - \hat{h})(x) > 0$ for all $x \in \mathscr{U}_\varepsilon$, that is $h_\varepsilon > h'_\varepsilon$ on $\mathscr{U}_\varepsilon$, contradicting the fact that $\int h_\varepsilon dm = \int h'_\varepsilon dm = 1$. Therefore $h_\varepsilon = h'_\varepsilon$ on $\mathscr{U}_\varepsilon$.
(ii) Suppose $\tilde{\Lambda}_0$ is not a least element; then $\tilde{\Lambda}_0 \to \tilde{\Lambda}$, where $\tilde{\Lambda}$ is a least element. Let h'_ε be the density of any acsm, i.e $\mathscr{L}_\varepsilon h'_\varepsilon = h'_\varepsilon$. We will show that $h'_\varepsilon = 0$ on $\tilde{\Lambda}_0$. We proceed by contradiction. First, since $\tilde{\Lambda}_0 \to \tilde{\Lambda}$, there is an ε-pseudo-orbit which starts from $\tilde{\Lambda}_0$ and ends up in $\tilde{\Lambda}$. Let h_ε denote the density of the unique acsm supported on $\tilde{\Lambda}$. If $h'_\varepsilon \neq 0$, we can invoke again the arguments of Lemma 1 and point (i) of this theorem to conclude that $h_\varepsilon > 0$ on $\tilde{\Lambda}$ too and also that $\min(h_\varepsilon, h'_\varepsilon)$ is a fixed

---

[7] The ergodicity insures this possibility for points in the same ergodic component; the equivalence relation allows to pass from one representative to the other in the least element, and finally the recursive construction of $\mathscr{U}_\varepsilon$ allows to get the external points $\mathscr{U}_\varepsilon/\Lambda$.

point of $\mathscr{L}_\varepsilon$. But the support of such a minimum is a subset of $\tilde{\Lambda}$ since for ε small enough, $h_\varepsilon = 0$ outside $\tilde{\Lambda}$. Then by uniqueness of the density h_ε over $\tilde{\Lambda}$ we get that $h_\varepsilon = \min(h_\varepsilon, h'_\varepsilon)$. This implies that $\int_M h'_\varepsilon dm > 1$, which is false.

Proof (of Corollary 3). By Proposition 1 and part (ii) of Theorem 1, for ε small enough, the support of any ergodic acsm μ_1^ε, intersects the support of a least element. By part (i) of Theorem 1 a small neighborhood of this least element supports an ergodic acsm μ_2^ε. By repeating the arguments of the proof of part (ii) of Theorem 1 we obtain that those two ergodic acsm must coincide.

4 Pseudo-orbits and metastability

In this section we assume that:
(M1) As an operator on $\mathscr{B}(M)$, $\mathscr{L}$ has 1 as an eigenvalue of multiplicity two. Moreover, if $\lambda_i \neq 1$ is an eigenvalue of $\mathscr{L}$, then $|\lambda| < 1$.
(M2) The map T has a unique *least element*.
Under conditions (M1) and (M2), we will show that random perturbations of T exhibit a *metastable behavior*[8]. For this purpose, we first introduce some notation and recall the Keller-Liverani perturbation theorem [12]. We adapt it to our situation which deals with the two adapted norms $||\cdot||_{\mathscr{B}} = |\cdot| + ||\cdot||_1$ and $||\cdot||_1$. For the unperturbed Perron-Frobenius operator $\mathscr{L}$ let us consider the set

$$V_{\delta,r}(\mathscr{L}) = \{z \in C : |z| \leq r \text{ or } \mathrm{dist}(z, \sigma(\mathscr{L})) \leq \delta\},$$

where $\sigma(\mathscr{L})$ is the spectrum of $\mathscr{L}$ as an operator on $\mathscr{B}$. Further, for $\varepsilon \geq 0$, we define the following operator norm

$$|||\mathscr{L}_\varepsilon||| = \sup_{||f||_{\mathscr{B}} \leq 1} ||\mathscr{L}_\varepsilon f||_1. \tag{2}$$

Note that by (LY) and (RLY), the operators $\mathscr{L}$, $\mathscr{L}_\varepsilon$ satisfy all the conditions of [12]. Thus, we are ready to state and use the following important result of [12]:

Theorem 2. *[12] If* $\lim_{\varepsilon\to 0} |||\mathscr{L}_\varepsilon - \mathscr{L}||| = 0$ *then for sufficiently small* $\varepsilon > 0$, $\sigma(\mathscr{L}_\varepsilon) \subset V_{\delta,r}(\mathscr{L})$*. Moreover, in each connected component of* $V_{\delta,r}(\mathscr{L})$ *that does not contain* 0 *both* $\sigma(\mathscr{L})$ *and* $\sigma(\mathscr{L}_\varepsilon)$ *have the same multiplicity; i.e., the associated spectral projections have the same rank.*

Using Theorem 2, we show that our random system $(\mathscr{X}_\varepsilon^n)$ exhibits metastable behavior:

Proposition 2. *Suppose that*

---

[8] By a metastable behavior we mean that $\mathscr{L}_\varepsilon$, as an operator on $\mathscr{B}$, will have 1 as a *simple* eigenvalue and will have another real eigenvalue ξ_ε very close to 1. Moreover ξ_ε has the second largest modulus among eigenvalues of $\mathscr{L}_\varepsilon$. Such a ξ_ε determines the rate of mixing [4] of the random system $(\mathscr{X}_n^\varepsilon)$.

- *T satisfies assumptions (M1), (M2);*
- $\lim_{\varepsilon\to 0} |||\mathscr{L} - \mathscr{L}_\varepsilon||| = 0$.

Then, as an operator on $\mathscr{B}$, $\mathscr{L}_\varepsilon$ has 1 *as a simple eigenvalue. Moreover, $\mathscr{L}_\varepsilon$ has a real eigenvalue ξ_ε very close to* 1. *In particular, ξ_ε has the second largest modulus among eigenvalues of $\mathscr{L}_\varepsilon$.*

Proof. Assumption (M1) states that the spectrum of $\mathscr{L}$, as an operator on $\mathscr{B}$, satisfies the following: $\exists$ an $r \in (\eta, 1)$[9] and a $\delta > 0$ such that:

1. The eigenvalue 1 of $\mathscr{L}$ is of multiplicity two;
2. if $\lambda_i \neq 1$ is an eigenvalue of $\mathscr{L}$, then $\lambda_i \in B(0,r)$;
3. $B(0,r) \cap B(1,\delta) = \emptyset$.

Moreover, under assumptions (M2), using Theorem 1, the random map $(\mathscr{X}_\varepsilon^n)$ has exactly one ergodic acsm; i.e., as an operator on $\mathscr{B}$, $\mathscr{L}_\varepsilon$ has 1 as a *simple* eigenvalue. Consequently, by Theorem 2, for sufficiently small ε, the spectrum of $\mathscr{L}_\varepsilon$ satisfies the following:

1. $\mathscr{L}_\varepsilon$ has a real eigenvalue $\xi_\varepsilon < 1$, with $\xi_\varepsilon \in B(1,\delta)$;
2. if $\lambda_{i,\varepsilon} \notin \{1, \xi_\varepsilon\}$ is an eigenvalue of $\mathscr{L}_\varepsilon$, then $\lambda_{i,\varepsilon} \in B(0,r)$.

Remark 2. The condition $\lim_{\varepsilon\to 0} |||\mathscr{L} - \mathscr{L}_\varepsilon||| = 0$ of Proposition 2 can be checked in several cases. A general theorem is presented in Lemma 8 of [5] for piecewise expanding maps of the interval endowed with our pair of adapted spaces where the noise is represented by a convolution kernel. In the multidimensional case, using quasi-Hölder spaces, the proof is given in Proposition 4.3 of [1].

Remark 3. Proposition 2 does not, in general, hold when the number of ergodic T-acim is $l > 2$. This is due to the fact that
a) If l is odd, then $l-1$ is even. Therefore, the transfer operator $\mathscr{L}_\varepsilon$ may have $l-1$ complex eigenvalues of modulus one sitting in $B(1,\delta)$.
b) If l is even, then $l-2$ is even. Therefore, the transfer operator $\mathscr{L}_\varepsilon$ may have $l-2$ complex eigenvalues of modulus one sitting in $B(1,\delta)$.

5 Random Transformations

In sections 2, 3 and 4 we studied random perturbations of a dynamical system in the framework of general Markov processes. Nevertheless, it is often useful to deal with the case when the Markov process is generated by *random transformations* [15]. In

---

[9] By (LY) and (RLY), η is an upper bound on the essential spectral radius of $\mathscr{L}$ and the essential spectral radius of $\mathscr{L}_\varepsilon$.



Fig. 1 An example of a 1−dimensional map T with two least elements

this setting, we consider an i.i.d. stochastic process $(\omega_k)_{k\in N}$ with values in Ω_ε and with probability distribution θ_ε. We associate with each $\omega \in \Omega_\varepsilon$ a map $T_\omega : M \to M$ and we consider the random orbit starting from the point x and generated by the realization $\underline{\omega}_n = (\omega_1, \omega_2, \cdots, \omega_n)$, defined as : $T_{\underline{\omega}_n} := T_{\omega_n} \circ \cdots \circ T_{\omega_1}(x)$. This defines a Markov process $\mathscr{X}_\varepsilon$ with transition function

$$P(x,A) = \int_{\Omega_\varepsilon} \mathbf{1}_A(T_\omega(x))d\theta_\varepsilon(\omega),$$

where $A \in \mathscr{B}(M)$, $x \in M$ and $\mathbf{1}_A$ is the indicator function of a set A. The transition function induces an operator $\mathscr{U}_\varepsilon^*$ which acts on measures μ on $(I, \mathscr{B}(M))$ as:

$$\mathscr{U}_\varepsilon^* \mu(A) = \int_M \int_{\Omega_\varepsilon} \mathbf{1}_A(T_\omega(x))d\theta_\varepsilon(\omega)d\mu(x) = \int_I \mathscr{U}_\varepsilon \mathbf{1}_A(x)d\mu_\varepsilon(x),$$

where $\mathscr{U}_\varepsilon$ is the random evolution operator acting on L^∞_m functions g:

$$\mathscr{U}_\varepsilon g = \int_{\Omega_\varepsilon} g \circ T_\omega d\theta_\varepsilon(\omega). \tag{3}$$



Fig. 2 An example of a 1−dimensional map T with one least element

A measure μ_ε on $(M, \mathscr{B}(M))$ is called a $\mathscr{X}_\varepsilon$-stationary measure if and only if, for any $A \in \mathscr{B}(M)$,

$$\mathscr{U}_\varepsilon^* \mu_\varepsilon(A) = \mu_\varepsilon(A). \tag{4}$$

We are interested in studying $\mathscr{X}_\varepsilon$-acsm. By (3), one can define the transfer operator $\mathscr{L}_\varepsilon$ (Perron-Frobenius) acting on $L^1(M, \mathscr{B}(M), m)$ by:

$$(\mathscr{L}_\varepsilon f)(x) = \int_{\Omega_\varepsilon} \mathscr{L}_\omega f(x) d\theta_\varepsilon(\omega), \tag{5}$$

which satisfies the duality condition

$$\int_M g \mathscr{L}_\varepsilon f dm = \int_M \mathscr{U}_\varepsilon g f dm \tag{6}$$

where g is in some subset of L_m^∞ and $\mathscr{L}_\omega$ is the transfer operator associated with T_ω. It is well know that $\mu_\varepsilon := \rho_\varepsilon m$ is a $\mathscr{X}_\varepsilon$-acsm if and only if $\mathscr{L}_\varepsilon \rho_\varepsilon = \rho_\varepsilon$; i.e., ρ_ε is a $\mathscr{X}_\varepsilon$-invariant density. In order to use the results of sections 2, 3 and 4, we also assume that assumptions (P1), (P2)[10] (B1)-(B3), (LY) and (RLY) hold. Moreover, we assume that (5) reduces to

$$(\mathscr{L}_\varepsilon f)(x) = \int_{\Omega_\varepsilon} \mathscr{L}_\omega f(x) d\theta_\varepsilon(\omega) = \int_M p_\varepsilon(z,x) f(z) dz. \tag{7}$$

In fact, an important example of a random perturbation where $\mathscr{L}_\varepsilon$ can be reduced as in (7) is the case of *additive* noise. For instance if $M = S^q$, the q-dimensional torus [11], let define $T_\omega = T(x) - \omega \bmod S^q$, where $\omega \in S^q$. Let the density of θ_ε with respect to the Lebesgue measure $d\omega$ on S^m, h_ε, be continuously differentiable density with support contained in the square $\Omega_\varepsilon \equiv [-\varepsilon, \varepsilon]^q$: $\int d\theta_\varepsilon = \int h_\varepsilon(\omega) d\omega = 1$. It is then straightforward to check that $p_\varepsilon(x,y) = h_\varepsilon(Tx - y)$.

5.1 1-*dimensional examples*

To illustrate our results, we present two examples of 1-dimensional maps[12]. In Example 1 we present a map that has three ergodic components and two least elements. In Example 2 we present a map which has two ergodic components and exactly one least element. Moreover, we show that when T of Example 2 is perturbed by an additive noise, it exhibits a metastable behavior.

Example 1. In this example $T : [0,10] \to [0,10]$. The graph of T is shown in Fig. 1. T is piecewise linear and Markov with respect to the partition:

$$[0,1), \dots, [4,5), [5,5.5) \dots, [9.5,10].$$

One can easily check that that T has exactly three ergodic acim whose supports $\Lambda_1, \Lambda_2, \Lambda_3$ are, respectively, equal to $[1,4]$, $[5.5,7.5]$, and $[7.5,9.5]$. Moreover, one can easily check that T admits two least elements. Namely, $\{\Lambda_1\}$ and $\{\Lambda_2, \Lambda_3\}$.

Example 2. In this example $T : [0,10] \to [0,10]$. The graph of T is shown in Fig. 2. T is piecewise linear and Markov with respect to the partition:

$$[0,1), \dots, [9,10].$$

One can easily check that that T has exactly two ergodic acim whose supports Λ_1, Λ_2 are, respectively, equal to $[1,5]$, and $[5,9]$. Moreover, one can easily check that T

---

[10] (P1) and (P2) in this setting are analogous to:
(P1) For all $x \in I$ the measure $P(x,\cdot)$ defined above on the Borel subsets of I by $P(x,A) = \theta_\varepsilon\{\omega \in \Omega_\varepsilon\ ,\ T_\omega(x) \in A\}$ is absolutely continuous with respect to Lebesgue, namely we have a summable density $p_\varepsilon(x,\cdot)$ such that: $P(x,A) = \int_A p_\varepsilon(x,y) dy$;
(P2) We have: support of $P(x,\cdot)$ coincides with $B_\varepsilon(Tx), \forall x \in I$.

[11] If M is not the torus we assume that, for all $\omega \in \Omega_\varepsilon$, $T_\omega(M) \subseteq M$.

[12] In both examples the Banach space $\mathscr{B}$ is considered to be the space of functions of bounded variation.

admits exactly one least elements. Namely, $\{\Lambda_1, \Lambda_2\}$. When T is perturbed by an additive noise, by using Lemma 8 of [5], we obtain that $\lim_{\varepsilon\to 0} |||\mathscr{L}_\varepsilon - \mathscr{L}||| = 0$. Consequently, by Proposition 2, the random system exhibits a metastable behavior.

Acknowledgements H. Hu thanks the University of Toulon and CPT Marseille where parts of this work was conducted. WB would like to thank the London Mathematical Society for supporting his visit to CPT Marseille where parts of this research was conducted. SV thanks the French ANR Perturbations, the CNRS-PEPS "Mathematical Methods for Climate Models" and the CNRS Pics "Propriétés statistiques des systémes déterministes et aléatoires" N. 05968, for support. SV thanks Ph. Marie for useful discussions.

References

1. Aytac, H., Freitas, J., Vaienti, S. *Laws of rare events for deterministic and random dynamical systems*. Submitted. Avalaible at arxiv.org/pdf/1207.5188
2. Bahsoun, W. , Vaienti, S. *Metastability of certain intermittent maps*. Nonlinearity, 25 (2012), no. 1, 107–124.
3. Bahsoun, W, Vaienti, S. *Escape rates formulae and metastability for randomly perturbed maps*. Submitted. Available at arXiv:1206.3654.
4. Baladi, V. *Positive transfer operators and decay of correlations*. Advanced Series in Nonlinear Dynamics, 16. World Sci. Publ., NJ, 2000.
5. Baladi, V., Young, L.-S. *On the spectra of randomly perturbed expanding maps*. Comm. Math. Phys. 156 (1993), no. 2, 355–385.
6. Dellnitz, M., Froyland, G., Horenkamp, C., Padberg-Gehle, K., and Sen Gupta, A. *Seasonal variability of the subpolar gyres in the Southern Ocean: a numerical investigation based on transfer operators*. Nonlinear Processes in Geophysics, 16:655-664, (2009).
7. Dolgopyat, D. and Wright, P. *The Diffusion Coefficient for Piecewise Expanding Maps of the Interval with Metastable States.* Stochastics and Dynamics, 12 (2012), no. 1, 1150005, 13 pp.
8. Froyland, G. and Stancevic, O. *Metastability, Lyapunov exponents, escape rates, and topological entropy in random dynamical systems*. To appear in Stochastics and Dynamics.
9. Gonzaléz-Tokman, C., Hunt, B., Wright, P. *Approximating invariant densities for metastable systems*, Ergod. Th. Dynam. Sys. (2011), 31, 1345–1361.
10. Hu, H., Vaienti, S. *Absolutely continuous invariant measures for non-uniformly expanding maps*. Ergod. Th. Dynam. Sys, 29 (2009), 1185-1215.
11. Keller, G. *Stochastic stability in some chaotic dynamical systems*. Monatsh. Math. 94 (1982), no. 4, 313–333.
12. Keller, G., Liverani, C. *Stability of the spectrum for transfer operators*. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 28 (1999), no. 1, 141-152.
13. Keller, G. and Liverani, C. *Rare events, escape rates, quasistationarity: some exact formulae*. J. Stat. Phys., 135 (2009), no. 3, 519-534.
14. Kifer, Y. *Attractors via random perturbations*, Commun. Math. Phys., (1989), 445–455.
15. Kifer, Y. *Ergodic theory of random transformations*, Birkhäuser, 1986.
16. Liverani, C. *Multidimensional expanding maps with singularities: a pedestrian approach*. Ergod. Th. Dynam. Sys. DOI:10.1017/S0143385711000939.
17. Ruelle, D. *Small random perturbations of dynamical systems and the definition of attractors*, Commun. Math. Phys., (1981), 82, 137–151.
18. Santitissadeekorn, N., Froyland, G., and Monahan, A. *Optimally coherent sets in geophysical flows: A new approach to delimiting the stratospheric polar vortex*. Phys. Rev. E, (2010), 056311-6.